



Fig. 17. World distribution of pipits, *Anthus*. See caption on p. 51.

- | | | | | |
|-------------------|---------------------|-------------------|------------------|-------------------|
| 1 - Short-tailed | 10 - Yellowish | 19 - Pechora | 28 - Grassveld | 37 - Plain-backed |
| 2 - Bushveld | 11 - Chaco | 20 - Rosy | 29 - African | 38 - Buffy |
| 3 - Sokoke | 12 - Ochre-breasted | 21 - Buff-bellied | 30 - Cameroon | 39 - Long-tailed |
| 4 - Correndera | 13 - Tree | 22 - Striped | 31 - Jackson's | 40 - Long-legged |
| 5 - South Georgia | 14 - Red-throated | 23 - African Rock | 32 - Long-billed | 41 - Tawny |
| 6 - Sprague's | 15 - Meadow | 24 - Mountain | 33 - Bannerman's | 42 - Berthelot's |
| 7 - Short-billed | 16 - Rock | 25 - Upland | 34 - Wood | 43 - Blyth's |
| 8 - Hellmayr's | 17 - Water | 26 - Richard's | 35 - Kimberley | 44 - Nilgiri |
| 9 - Paramo | 18 - Olive-backed | 27 - Paddyfield | 36 - Malindi | 45 - Alpine |